

43

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
8 November 2001 (08.11.2001)

PCT

(10) International Publication Number
WO 01/82962 A2

- (51) International Patent Classification⁷: A61K 39/12
- (21) International Application Number: PCT/CA01/00577
- (22) International Filing Date: 25 April 2001 (25.04.2001)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
60/200,011 27 April 2000 (27.04.2000) US
- (71) Applicant (*for all designated States except US*): AVEN-
TIS PASTEUR LIMITED [CA/CA]; 1755 Steeles Av-
enue West, Toronto, Ontario M2R 3T4 (CA).
- (72) Inventors; and
- (75) Inventors/Applicants (*for US only*): ROVINSKI, Ben-
jamin [CA/CA]; 70 Winding Lane, Thornhill, Ontario L4J
5H6 (CA). TARTAGLIA, James [US/US]; 7 Christina
Drive, Schenectady, NY 12303 (US). CAO, Shi-Xian
[CA/CA]; Apartment 408, 716 The West Mall, Etobicoke,
Ontario M9C 4X6 (CA). PERSSON, Roy [CA/CA]; Unit
604, 7 Bishop Avenue, Toronto, Ontario M2M 4J4 (CA).
KLEIN, Michel, H. [CA/CA]; 16 Munro Boulevard,
Toronto, Ontario M2P 2P9 (CA).
- (74) Agent: STEWART, Michael, I.; Sim & McBurney, 6th
Floor, 330 University Avenue, Toronto, Ontario M5G 1R7
(CA).
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM,
HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK,
LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX,
MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL,
TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- Published:
— without international search report and to be republished
upon receipt of that report
- For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*



WO 01/82962 A2

(54) Title: IMMUNIZING AGAINST HIV INFECTION

(57) Abstract: A virus neutralizing level of antibodies to a primary HIV isolate is generated in a host by a prime-boost administration of antigens. The primary antigen is a DNA molecule encoding an envelop glycoprotein of a primary isolate of HIV-1 while the boosting antigen is either a non-infectious, non-replicating HIV-like particle having the envelope glycoprotein of a primary isolate of HIV-1 or an attenuated viral vector expressing an envelope glycoprotein of a primary isolate of HIV-1.

TITLE OF INVENTION
IMMUNIZING AGAINST HIV INFECTION

FIELD OF THE INVENTION

[0001] The present invention relates to the field of immunology and, in particular, to methods and compositions for immunizing a host against infection with HIV.

BACKGROUND OF THE INVENTION

[0002] Human immunodeficiency virus is a human retrovirus and is the etiological agent of acquired immunodeficiency syndrome (AIDS). It is estimated that more than 33 million people have been infected with HIV world-wide as of December 1999 (Ref 1- various references are referred to in parenthesis to more fully describe the state of the art to which this invention pertains. Full bibliographic information for each citation is found at the end of the specification, immediately preceding the claims. The disclosure of these references are hereby incorporated by reference into the present disclosure).

[0003] As the HIV epidemic continues to spread world wide, the need for an effective vaccine remains urgent. Efforts to develop such a vaccine have been hampered by several factors three of which are: (a) the extraordinary ability of the virus to mutate; (b) inability of most known specificities of anti-HIV antibodies to neutralise HIV primary isolates consistently; and (c) lack of understanding of the correlates of protective immunity to HIV infection. Over the last 10 years, several candidate HIV vaccines have been tested in primates for their immunoprotective abilities (Ref. 2). These studies suggest that both neutralising antibodies and cell-mediated immunity play a role in conferring sterilizing immunity and preventing progression towards disease (Ref 3, 4). While the correlates for immune protection against HIV-1 infection are currently unknown, an effective HIV vaccine should elicit both strong neutralising antibody and cytotoxic T lymphocyte (CTL) responses.

[0004] Envelope subunit vaccines have been shown to induce high titred humoral responses, but were inefficient in eliciting CTL responses (Ref 5). Live recombinant pox vectors have been shown to elicit very potent CTL responses, however these vectors were ineffective for generating a significant antibody

response (Ref 6). In attempts to combine the two immunization types, several clinical trials involved a prime-boost strategy, consisting of initial viral vector immunization followed by boosts with recombinant HIV-1 envelope subunits (Ref 7, 8), have led to limited success with respect to CTL responses. Other vaccine
5 approaches have used non-infectious, non-replicating, immunogenic virus-like particles (VLP) for immunising against HIV infection (Ref 9, 10). This type of immunogen has lead to the generation of neutralizing antibodies to a laboratory HIV-1 strain (Ref 10).

[0005] A prime-boost approach has been investigated using non-infectious
10 VLPs to enhance HIV-specific CTL responses in mice primed with recombinant canarypox vector vCP205 encoding HIV-1gp 120 (MN strain) (Ref 11). This study showed that VLPs could boost the CTL response to the canarypox vector.

[0006] Recently, a study showing the induction of neutralizing antibodies to a HIV-1 primary isolate in chimpanzees has been reported (Ref 12). In this
15 study, recombinant adenovirus expressing gp160 was used as the priming agent and recombinant gp120 protein was used to boost the monkeys.

[0007] There is still a need for vaccines and immunization regimes to induce both a strong CTL response as well as neutralizing antibodies to HIV primary isolates.

20 SUMMARY OF THE INVENTION

[0008] In accordance with one aspect of the present invention, there is provided a method for generating, in a host, particularly a human host, a virus neutralizing level of antibodies to a primary HIV isolate, comprising at least one administration of a priming antigen to the host, wherein the priming antigen
25 comprises a DNA molecule encoding an envelope glycoprotein of a primary isolate of HIV, resting the host for at least one specific resting period to provide for clonal expansion of an HIV antigen specific population of precursor B-cells therein to provide a primed host, and at least one administration of a boosting antigen to the primed host to provide said neutralizing levels of antibodies,
30 wherein the boosting antigen is selected from the group consisting of a non-infectious, non-replicating, immunogenic HIV-like particle having at least part of the envelope glycoprotein of a primary isolate of HIV and an attenuated viral

vector expressing at least part of an envelope glycoprotein of a primary isolate of HIV.

[0009] The primary HIV isolate may be an HIV-1 isolate including from the clade B HIV-1 clinical isolate HIV-1_{Bx08}, although any other primary HIV-1 isolate may be employed in the immunization procedures of the invention.

[0010] The DNA molecule encoding the envelope glycoprotein of a primary isolate of HIV may be contained in a plasmid vector under the control of a heterologous promoter, preferably a cytomegalovirus promoter, for expression of the envelope glycoprotein in the host, which may be a human host.

[0011] The vector utilized for DNA molecule immunization is novel and constitutes a further aspect of the present invention. Preferably, the vector has the identifying characteristics of pCMV3Bx08 shown in Figure 2, such identifying characteristics being the nucleic acid segments and restriction sites identified in Figure 2.

[0012] A priming administration of antigen may be effected in a single or in multiple administrations of the priming antigen. In the latter case, the at least one specific resting period to permit clonal expression of HIV antigen-specific population precursor B-cells may be effected after each priming administration. The at least one specific resting period may be between about 2 and 12 about months.

[0013] In the embodiment where the boosting antigen is a non-infectious, non-replicating, immunogenic HIV-like particle, such particle may comprise an assembly of:

- (i) an *env* gene product,
- (ii) a *pol* gene product, and
- (iii) a *gag* gene product

with the particle being encoded by a modified HIV genome deficient in long terminal repeats (LTRs) and containing *gag*, *pol* and *env* in their natural genomic arrangement. Such particles and the manufacture thereof are described in US Patent No. 5,439,809, assigned to the assignee hereof and the disclosure of which is incorporated herein by reference. Such particles can include mutations in *gag* and *pol* to further reduce potential infectivity, as more fully described in United

States Patent No. 6,080,408, assigned to the assignee hereof and the disclosure of which is incorporated herein by reference (WO 96/06177). In a preferred embodiment, the *env* gene is that from primary isolate BX08. The *gag* gene and *pol* gene may be those from the same primary isolate or may be chosen from those of other HIV-1 isolates, which may be primary isolates.

[0014] The non-infectious, non-replicating, immunogenic HIV-like particle may be administered in conjunction with an adjuvant. Any suitable adjuvant may be used, such as QS21, DC-chol, RIBI or Alum.

[0015] Such non-infectious, non-replicating, immunogenic HIV particle may be formed by expression from a suitable vector in mammalian cells. In accordance with an additional aspect of this invention, there is provided a vector comprising a modified HIV-genome deficient in long terminal repeats and a heterologous promoter operatively connected to said genome for expression of said genome in mammalian cells to produce the non-infectious, non-replicating and immunogenic particle, wherein at least the *env* gene of the modified HIV-genome is that from a primary isolate of HIV. The *gag* and *pol* genes of the modified HIV genome may be those from the same primary isolate or those from another isolate, which may be a primary isolate.

[0016] The vector preferably is a plasmid vector while the primary isolate preferably is BX08. The promoter may be the metallothionein promoter. The vector preferably has the identifying characteristics of plasmid p133B1 shown in Figure 3, such identifying characteristics being the nucleotide segments and restriction sites identified in Figure 3.

[0017] In the embodiment where the boosting antigen is an attenuated viral vector, the attenuated viral vector may be an attenuated avipox virus vector, particularly the attenuated canary poxvirus ALVAC. The attenuated viral vectors used herein form another aspect of the invention. The attenuated viral vector may contain a modified HIV genome deficient in long terminal repeats (LTRs), wherein at least the *env* gene is that from primary isolate BX08. The *gag* and *pol* genes of the modified genome may be those from the same primary isolate or may be chosen from other HIV isolate.

[0018] The attenuated canarypox virus-based vector ALVAC is a plaque-cloned derivative of the licensed canarypox vaccine, Kanapox, and is described in reference 19. The attenuated canary pox vector preferably has the identifying characteristics of vCP1579 shown in Figure 4, such identifying characteristics
5 being the nucleic acid segments and restriction sites identified in Figure 4.

[0019] The at least one administration of a boosting antigen may be effected in a single administration or at least two administration of the boosting antigen.

[0020] The invention further includes compositions comprising the
10 immunogens as provided herein and their use in the manufacture and formulation of immunogenic compositions including vaccines.

BRIEF DESCRIPTION OF THE DRAWINGS

[0021] The present invention will be further understood from the following description with reference to the drawings, in which:

15 Figure 1 shows the details of the elements of plasmid pCMVgDtat⁻vpr⁻Bx08.

Figure 2 shows the details of the elements of plasmid pCMV3Bx08.

Figure 3 shows the details of the elements of plasmid p133B1.

20 Figure 4 shows the details of the insertions into ALVAC (2) to provide vector vCP1579.

Figures 5A and 5B contain a representation in time-line form of the immunization regime used wherein the study groups are described in Table 1. The numbers below the lines refer to weeks.

25 Figure 6 shows the immunoreactivity to HIV-1 antigens of the serum diluted 1:100 from the macaques immunized with the various preparations as described in Table 1.

Figure 7 shows the immunoreactivity to HIV-1 antigens of the serum diluted 1:1000 from the macaques immunized with the various preparations
30 as described in Table 1.

Figure 8 shows the details of the elements of pMPC6H6K3E3.

Figure 9 shows the details of the elements of pMPC5H6PN.

Figure 10 shows the details of the elements of pHIV76.

Figure 11 shows the nucleotide sequence (SEQ ID NO: 1) for the H6/HIV Pol/Nef epitope cassette in the ALVAC C5 site of vCP1579.

Figure 12 contains the nucleotide sequence of C6 region (coding strand SEQ ID NO: 16, complementary strand SEQ ID NO: 17, K3L amino acid sequence SEQ ID NO: 18, E3L amino acid sequence SEQ ID NO: 19).

GENERAL DESCRIPTION OF INVENTION

[0022] As noted earlier, the present invention involves administration of HIV antigens to elicit virus-neutralizing levels of antibodies against a primary HIV isolate.

[0023] A DNA construct was prepared incorporating the *env* gene from the primary isolate Bx08 under the control of the cytomegalovirus promoter and the construct, pCMV3Bx08, is shown in Figure 2. The construct pCMV3Bx08 is derived from plasmid pCMVgDtat vpr Bx08 seen in Figure 1. The DNA construct pCMV3Bx08 was used in a priming immunization step to a host, macaque monkeys being the animal model chosen.

[0024] Following the priming immunization step, which may be effected in one or more administrations of the DNA construct, the host is allowed to rest to provide for clonal expression of an HIV antigen specific population of precursor B-cells therein to provide a primed host.

[0025] The boosting administration is effected either with a non-infectious, non-replicating, immunogenic HIV-like particle (VLP) or an attenuated viral vector.

[0026] For this purpose, a VLP expression plasmid was constructed containing a modified HIV genome lacking long terminal repeats in which the *env* gene is derived from primary isolate BX08, wherein the modified HIV genome is under the control of a metallothionein promoter. The construct, p133B1, shown in Figure 3, was used to effect expression in mammalian cells of the non-infectious, non-replicating, immunogenic HIV-like particles, in which the *env* gene product is that from the primary isolate BX08.

[0027] In the case of the attenuated virus vector, a recombinant attenuated canarypox virus vector was constructed to contain the *env* gene from primary

isolate BX08. The viral vector vCP1579 (Figure 4) was prepared by a variety of manipulations from plasmid pHIV76 (Figure 10), as shown described in detail below.

[0028] These products were utilized in a boosting administration to the primed macaques. The boosting administration may be effected in one or more immunizations. In a preferred aspect of the invention, the non-infectious, non-replicating immunogenic HIV-like particles may co-administered with the DNA construct in the priming administration and the DNA construct may be coadministered with the HIV-like particles in the boosting administration.

[0029] Immunizations were effected in accordance with the procedure of the invention and the results obtained were compared with those obtained using other protocols according to the protocols set forth in Table 1. The immunization regimes used are shown as time lines in Figures 5A and 5B.

[0030] The results obtained following the various protocols showed that, in particular, a primary DNA vaccination in combination with a boost from either the VLP or the attenuated canarypox virus enhanced the levels of neutralizing antibodies, as indicated by the reduction of detectable p24 levels in cells infected with primary HIV isolates.

Biological Deposits

[0031] Certain vectors that are described and referred to herein have been deposited with the American Type Culture Collection (ATCC) located at 10801 University Boulevard Manassas, Virginia 20110-2209, USA, pursuant the Budapest Treaty and prior to the filing of this application. Samples of the deposited vectors will become available to the public and all restrictions imposed or access to the deposits will be received upon grant of a patent based on this United States patent application or the United States patent application in which they are described. In addition, the deposit will be replaced if viable samples cannot be dispensed by the Depository. The invention described and claimed herein is not limited in scope by the biological materials deposited, since the deposited embodiment is intended only as an illustration of the invention. Any equivalent of similar vectors that contain nucleic acids which encode equivalent or

similar antigens as described in this application are within the scope of the invention.

Deposit Summary

| | <u>Plasmid</u> | <u>ATCC</u> | <u>Deposit Date</u> |
|---|----------------|-------------|---------------------|
| 5 | pMT-HIV | 40912 | October 12, 1990 |
| | pCMVgDtat-vpr | 209446 | November 11, 1997 |

EXAMPLES

[0032] The above disclosure generally describes the present invention. A more complete understanding can be obtained by reference to the following specific Examples. These Examples are described solely for purposes of illustration and are not intended to limit the scope of the invention. Changes in form and substitution of equivalents are contemplated as circumstances may suggest or render expedient. Although specific terms have been employed herein, such terms are intended in a descriptive sense and not for purposes of limitation.

15 [0033] Example 1

[0034] This Example describes the construction of plasmid pCMV3BX08.

[0035] The plasmid, pCMV3BX08, contains sequence segments from various sources and the elements of construction are depicted in Figure 2.

[0036] The prokaryotic vector pBluescript SK (Stratagene) is the backbone of the plasmid pCMV3.BX08 and was modified by the replacement of the *Amp*^R with *Kan*^R gene and the deletion of the *fl* and the *LacZ* region. To achieve the desired modifications, the sequence between *AhdI* (nucleotide 2,041) and *SacI* (nucleotide 759) of pBluescript SK, which contains the *Amp*^R, *fl* origin and the *LacZ*, was deleted. A 1.2 kb *PstI* fragment from the plasmid pUC-4K (Pharmacia) containing the *Kan*^R gene, was blunt end ligated to the *AhdI* site of pBluescript SK in a counter-clockwise orientation relative to its transcription. A 1.6 kb *SspI/PstI* DNA fragment containing the human cytomegalovirus immediate-early gene promoter, enhancer and intron A sequences (CMV) was ligated to the other end of the *Kan*^R gene so that the transcription from the CMV promoter proceeds in the clockwise orientation. A synthetic oligonucleotide segment containing translation initiation sequence and sequences encoding the human tissue plasminogen activator signal peptide (TPA) was used to link the

CMV promotor and the sequences encoding the envelope gene of the primary isolate HIV-1_{BX08}.

[0037] The envelope gene from the HIV-1 primary isolate BX08 was isolated from the plasmid pCMVgDtat⁻vpr⁻Bx08 illustrated in Figure 1. The
5 plasmid pCMVgDtat⁻vpr⁻Bx08 was derived from the deposited plasmid pCMVgDtat⁻vpr⁻, the construction of which is described in copending United States Patent Application No. 08/991,773 filed December 16, 1997, assigned to the assignee hereof and the disclosure of which is incorporated herein by reference, (WO 99/31250). The plasmid pCMVgDtat⁻vpr⁻Bx08 was derived by
10 substituting the BX08 envelope sequence from clade B HIV-1 clinical isolate HIV-1_{BX08} for the modified HIV genome sequence present in pCMVgDtat⁻vpr⁻. Plasmid pCMVgDtat⁻vpr⁻Bx08 was restricted with the restriction enzyme *Xho* I and made blunt ended with Klenow treatment. A *Not* I partial digestion was then performed and the resulting 6.3 kb fragment containing the *env* gene was isolated.
15 Plasmid pCMV3 (Invitrogen) was restricted with *Bam* HI and made blunt ended with Klenow treatment. The plasmid pCMV3 was then restricted with *Not* I and the resulting 4.4 kb fragment was isolated. The 6.3 and 4.4 kb fragments were ligated together to produce plasmid pCMV3BX08 (Figure 2).

[0038] The pCMV3BX08 construct was introduced into HB101 competent
20 cells according to manufacturer's recommendations (GibcoBRL). Correct molecular clones were identified by restriction and sequencing analysis and their expression of envelope glycoprotein was examined in transient transfections followed by Western blot analysis.

[0039] All DNAs used for immunizations were prepared using EndoFree
25 Plasmid Kit (Qiagen). For intramuscular immunizations either 3 mg or 600 µg of pCMVBX08, in 100 µl PBS was injected.

[0040] Proviral DNA for clade B HIV-1 clinical isolate HIV-1_{BX08} originated at Transgene (Strasbourg, France) and was isolated from genomic DNA of cells infected with the virus.

30 [0041] Example 2

[0042] This Example describes the construction of plasmid p133B1.

[0043] A Bx08 plasmid expression vector (p133B1, Figure 3) used to transfect the mammalian cells was engineered in several stages using pUC18 as the initial host plasmid. First, an 8.3-kbp fragment of HIV-1_{LAI} provirus encoding the gag, pol and env proteins was isolated. This fragment lacked the transcription regulatory elements and long terminal repeat elements from each end of the proviral genome to ensure the virus-like particles would be replication-incompetent. This fragment was linked to an inducible human type IIA metallothionein (MTIIA) promoter (Ref 13) and also to a Simian Virus 40 polyadenylation (polyA) addition/transcription termination sequence from plasmid pSV2dhfr (Ref 14). The modified fragment was then inserted into the pUC18 host vector. The resulting expression construct, named pMT-HIV, was used to transfect into African green monkey kidney (Vero) and COS monkey kidney cells. The procedure for obtaining pMT-HIV is further described in the aforementioned US Patent No. 5,439,809. Both transfected cell lines produced non-replicating virus-like particles when induced with metal ions (Ref 15).

[0044] Two further modifications were made to the proviral DNA in pMT-HIV to provide additional safety features to protect human cells against recombination events with reverse-transcribed DNA:

- 1) inactivation of the RNA packaging sequences; and
- 2) deletion of a large section of the *pol* gene encoding reverse transcriptase and integrase.

[0045] To delete the first RNA packaging signal, part of the DNA corresponding to the untranslated leader sequence of the mRNA was replaced with synthetic DNA lacking a 25-bp motif corresponding to nucleotides 753-777 (the *psi* sequence). To inactivate the second RNA packaging signal, two adenosine residues within a *gag* gene zinc finger sequence were changed to thymidine residues. Each of these residue changes had the effect of replacing cysteine residues in a Cys-His array with a serine in the gene product.

[0046] The *pol* gene deletion was effected by replacing a 1.9-kbp fragment with synthetic DNA containing stop codons in all three reading frames. This prevented read-through translation of the residual integrase coding sequence on the 3' side of the deletion. The 1.9-kbp deletion in *pol* also eliminated the

expression of reverse transcriptase and integrase enzymes. However, the deletion left intact the gene encoding the viral protease, which is both an immunogenic component of HIV-1 virus particles and allows the expression of particles with processed gag antigens closely resembling native virions (Ref 16). The protease
5 also contains epitopes that are conserved across HIV-1 clades. The modifications described with respect to *gag* and *pol* genes are more fully described in the aforementioned United States Patent No. 6,080,408 (WO 96/06177).

[0047] Finally, the HIV-1_{LAI} *env* gene within pMT-HIV was replaced with that of HIV-1_{Bx08}. To effect this replacement, a 2440-bp fragment containing the
10 *env* gene of Bx08 was amplified by polymerase chain reaction (PCR) from cells infected with this isolate. The PCR product was then used to replace the corresponding region present in pMT-HIV. However, the incoming fragment from HIV-1_{Bx08} was 125-bp shorter than the original HIV-1_{LAI} region owing to a deletion in the untranslated region between the *env* gene stop codon and the
15 termination/polyA addition sequence. The resulting construct replaced all but eleven amino acid residues of the LAI envelope proteins gp120 and gp41. Of these eleven, only the first three differ between the LAI and Bx08 isolates, and these are all charge-conservative changes meaning the final expression vector (p133B1) encoded a nearly authentic HIV-1_{Bx08} *env* protein.

20 [0048] Example 3

[0049] This Example describes the production of HIV-like particles.

[0050] African green monkey kidney (Vero) cells were recovered and cultivated in Dulbecco's modified Eagle medium (DMEM) containing 10% v/v fetal bovine serum (FBS), referred to below as Complete Medium. At passage
25 141, the cells were transfected with p133B1 using the calcium phosphate method when at approximately 30% confluence. The cells were shocked with glycerol 8 hours after transfection. For this step, six 10-cm dishes containing approximately 3.0×10^6 cells each in 10.0 mL of Complete Medium were prepared. Each dish received 25.0 μ g of expression vector and 2.0 μ g of plasmid pSV2neo (Ref 17).
30 The pSV2neo contains a selectable marker gene conferring resistance to the antibiotic geneticin (G418). Two days after transfection, the cells from each dish

were recovered by trypsinization and replated into twenty-five fresh dishes in Complete Medium supplemented with 0.5 mg/mL of G418.

[0051] In total, 394 colonies were isolated from the dishes using cloning cylinders. Each colony was recovered by trypsinization and divided into two cluster dish wells, one of the wells per clone was induced after reaching 50% to 90% confluence. Prior to induction, the wells were treated by replacing all the medium with fresh Complete Medium containing 10.0 μ M 5-azacytidine. After incubating for between 18 hours and 22 hours, the medium was removed and replaced with fresh DMEM containing 0.2% v/v FBS, 2.0 μ M CdCl₂ and 200.0 μ M ZnCl₂. The wells were incubated for a further 20 hours to 24 hours at which time samples of the medium were removed and tested by p24 ELISA.

[0052] The twenty highest-producing clones, based on the p24 titre, were chosen and cells from the corresponding uninduced wells were sub-cultured into one T-25 and one T-150 flask per clone. Both flasks were grown to confluence. The cells from the T-150 were recovered by trypsinization and cryopreserved at passage number 145. The cells from the T-25 were recovered by trypsinization every 3 days to 4 days and maintained up to passage 153. The cells were induced as above and samples retested by p24 ELISA at two different passages prior to passage 153.

[0053] The two highest p24 producers were chosen and were recovered by trypsinization every 3 days to 4 days up to passage 163. Samples from the clones were tested by p24 and gp120 ELISA from passage 158 and by p24 ELISA at passage 163, to assess clonal stability. The most suitable of these two cell lines, named 148 to 391, was chosen for further sub-cloning. The clone nomenclature defines the experiment number for this procedure, which was 148, and the number of the clone, which was number 391 of the original 394 isolated.

[0054] The vero cells were grown for approximately 100 h to 103 h and the medium was then replaced with growth medium containing 5-azacytidine. The bottles were then incubated for a further 20 h to 22 h, at which time the medium was replaced with serum-free medium containing CdCl₂ and ZnCl₂. The bottles were then incubated for 29 h to 31 h, at which time the medium was harvested, pooled and stored at 2°C to 8°C prior to purification.

[0055] The next day after harvesting, the solution was clarified, concentrated and diafiltered against phosphate buffer. The concentrate was passed through a ceramic hydroxyapatite (type I) column and the run-through was collected. The run-through from two successive sublots was pooled together and
5 pumped onto a sucrose density gradient in a continuous zonal ultracentrifuge rotor. Pseudovirion-containing fractions were collected and pooled. The pooled pseudovirion fractions were diafiltered against PBS containing 2.5% sucrose to reduce the sucrose content, concentrated and diafiltered again. The material was sterile filtered using a 0.2 μ m filter. At this stage the materials was designated as
10 a purified sub-lot and were stored at 2 to 8°C.

[0056] The adjuvants were prepared separately and filter sterilized before filling in single dose vials. QS21 was stored at -20°C.

[0057] Example 4

[0058] This Example describes the production of recombinant pox virus
15 vCP1579.

[0059] Recombinant pox virus vCP1579 (Figure 4) contains the HIV-1 gag and protease genes derived from the HIV-1 IIIB isolate, the gp120 envelope sequences derived from the HIV-1 Bx08 isolate, and sequences encoding a polypeptide encompassing the known human CTL epitopes from HIV-1 Nef and
20 Pol.

[0060] Recombinant vCP1579 (Figure 4) was generated by insertion of the vector modifying sequences from pMPC6H6K3E3 (Figure 8) encoding E3L and K3L into the C6 site of recombinant vCP1566 (Figure 4). Recombinant vCP1566 was generated by insertion of an expression cassette encoding a synthetic
25 polypeptide containing Pol CTL epitopes and Nef CTL epitopes (Figure 11) and plasmid pMPC5H6PN (Figure 9) into vCP1453 at the insertion site known as C5. Recombinant vCP1453 was generated by co-insertion of genes encoding HIV-1 env and gag/protease gene products, plasmid pHIV76 (Figure 10), into the ALVAC genome at the insertion site known as C3.

30 [0061] The construction of recombinant pox vectors containing the E3L and K3L genes has been described in United States patent 6,004,777 issued Dec 21, 1999 to Tartaglia et al. and the recombinant pox vectors describing the

insertion of HIV genes has been described in United States patent 5,766,598 issued June 16 1998 to Paoletti et al.

- [0062] The locus designated C3 was used for the insertion of the HIV-1 *env* and *gag* gene sequences into the ALVAC(2) vector, and the locus designated as C5 was the insertion site for the sequences encoding the HIV-1 Nef and Pol CTL epitopes. By virtue of the C3 and C5 loci existing within the extensive inverted terminal repetitions (ITRs) of the virus genome (approximately 41 kbp), insertion into these loci results in the occurrence of two copies of the inserted HIV-1 sequences.
- 10 [0063] Briefly, expression cassette pHIV76 (Figure 10) was engineered in the following manner. Plasmid p133B1 (Figure 3) containing the HIV-1Bx08 gp160 gene was used as the starting plasmid. The 3'-end of the H6 promoter was cloned upstream of the gp160 gene and three poxvirus early transcription termination signal sequences (T₅NT) were modified. This was accomplished by
- 15 cloning a 2,600 bp *Bam*HI-digested PCR fragment, containing the 3'-end of the H6 promoter and the T₅NT-modified HIV-1 (BX08) gp160 gene, into the *Bam*HI site of pBS-SK. This PCR fragment was generated from four overlapping PCR fragments (a 570 bp fragment, a 140 bp fragment, a 500 bp fragment and a 1,450 bp fragment) and the oligonucleotides, RW835 (5'-ATCATCATCGGATCC
- 20 CGGGGTCGCGATATCCGTTAAGTTTGTATCGTAATGAAAGTGAAGGAC C-3' - SEQ ID NO: 2) and RW836 (5'-ATCATCATCGGATCCCGGGGTT ATAGCAAAGCCCTTTC-3' - SEQ ID NO: 3). The 570 bp PCR fragment, containing the 3'-end of the H6 promoter and the 5'-end of the gp160 gene, was generated from the plasmid, p133B1, with the oligonucleotides, RW835 (5'-ATC
- 25 ATCATCGGATCCCGGGGTCGCGATATCCGTTAAGTTTGTATCGTAATG AAAGTGAAGGAGACC-3') and RW868 (5'-ATCAAGACTATAGAAGA GTGCATATTCTCTCTTCATC-3'). The 140 bp PCR fragment, containing an interior portion of the gp160 gene, was generated from plasmid p133-B1 with the oligonucleotides, RW864 (5'-GCACTCTTCTATAGTCTTGATATAGTAC-3' -
- 30 SEQ ID NO: 4) and RW865 (5'-AGCCGGGGCGCAGAAATGTATG GGAATTGGCAC-3' - SEQ ID NO: 5). The 500 bp PCR fragment, containing an interior portion of the gp160 gene, was generated from 133-3 with the

oligonucleotides, RW866 (5'-ATACATTTCTGCGCCCCGGCTGGT
TTTGCGATTC-3' - SEQ ID NO: 6) and RW867 (5'-GAAGAATTC
CCCTCCACAATTA AAC-3' - SEQ ID NO: 7). The 1,450 bp PCR fragment,
containing the 3'-end of the gp160 gene, was generated from p133-B1 with the
5 oligonucleotides, RW869 (5'-TGTGGAGGGGAATTCTTCTACTGTAATAC
AACACAAC-3' - SEQ ID NO: 8) and RW836 (5'-ATCATCATCGGAT
CCCGGGGTTATAGCAAAGCCCTTTC-3' - SEQ ID NO: 9). The 3'-end of the
570 bp PCR fragment overlaps the 5'-end of the 140 bp PCR fragment. The 3'-
end of the 140 bp PCR fragment overlaps the 5'-end of the 500 bp PCR fragment.
10 The 3'-end of the 500 bp PCR fragment overlaps the 5'-end of the 1450 bp PCR
fragment. The plasmid generated by this manipulation is called pRW997.

[0064] The sequence encoding gp41 was then replaced with the sequence
encoding the gp160 transmembrane (TM) region. This modification was
accomplished by cloning a 200 bp *MfeI-HindIII*-digested PCR fragment,
15 containing the 3'-end of the gp120 gene and the TM sequence, into the 4,400 bp
MfeI-HindIII fragment of pRW997. This PCR fragment was generated from two
overlapping PCR fragments (a 170 bp fragment and a 125 bp fragment) with the
oligonucleotides, HIVP97 (5'-TAGTGGGAAAGAGATCTTCAGACC-3' - SEQ
ID NO: 10) and HIVP101 (5'-TTTTAAGCTTTTATCCCTGCCTAACT
20 CTATTCAC TAT-3' - SEQ ID NO: 11). The 170 bp PCR fragment was
generated from pRW997 with the oligonucleotides, HIVP97 (5'-
TAGTGGGAAAGAGATCTTCAGACC-3' - SEQ ID NO: 12) and HIVP100 (5'-
CCTCCTACTATCATTATGAATATTCTTTTTTCTCTCTGCACCACTCT-3' -
SEQ ID NO: 13). The 125 bp PCR fragment was generated from pRW997 with
25 the oligonucleotides, HIVP99 (5'-AGAGTGGTGCAGAGAGAAAAA
AGAATATTCATAATGATAGTAGGAGGC-3' - SEQ ID NO: 14) and HIVP101
(5'-TTTTAAGCTTTTA TCCCTGCCTAACTCTATTCAT-3' - SEQ ID NO:
15). The plasmid generated by this manipulation is called pHIV71.

[0065] The H6-promoted gp120+TM gene was then cloned between C3
30 flanking arms, into a plasmid containing the I3L-promoted HIV1 gag/(pro) gene.
This modification was accomplished by cloning the 1,600 bp *NruI-XhoI* fragment
of pHIV71, containing the H6-promoted gp120+TM gene, into the 8,200 bp *NruI*-

*Xho*I fragment of pHIV63 . The plasmid generated by this manipulation is called pHIV76 (Figure 10). Plasmid pHIV76 was used in *in vivo* recombination experiments with ALVAC (CPpp) as rescue virus to yield vCP1453.

[0066] The sequence of the *nef/pol* regions is shown in Figure 12 and the
5 E3L and K3L sequences are shown in Figure 13. To generate ALVAC(2)120(BX08)GNP (vCP1579), expression cassettes consisting of the promoter/HIV-1 gene combinations were subcloned into an ALVAC donor plasmid, which were then used to insert the expression cassettes into defined sites in the ALVAC genome by *in vitro* recombination as previously described (Ref
10 20).

[0067] Example 5

[0068] This Example describes the results of immunization regimes.

[0069] Groups of four animals (macaques) each were randomly assigned to seven vaccine groups as illustrated in Table 1. In this Table, "BX08 DNA" refers to pCMV3BX08, prepared as described in Example 1, "BX08 VLP" refers to the pseudovirions produced by expression vector p133B1 in Vero cells, as described in Example 3, and "ALVAC(2) BX08" refers vCP1579, prepared as described in Example 4. Reference (pre-bleed) sera were sampled at -6 and -2 weeks pre-vaccination. Primary immunizations with the various vaccines were
15 given on weeks 0 and 4 with boosts on weeks 24 and 44 (Figures 5A, 5B). The vaccines were immunized intramuscularly into one quadricep of each macaque monkey.

[0070] Sera were prepared from whole-blood using SST collection tubes and analyzed using commercially available HIV-1 western blots. Groups 1, 2 and
25 7 showed low levels of anti-Env antibodies after the first boost (Figures 6 and 7). Based on ELISA values, the anti-env antibody levels were below 1 µg/ml of specific IgG. High levels of anti-gag antibodies were detected in groups 1, 2, 3, 4, and 7 (Figures 6 and 7). No HIV-1 specific antibodies were detected in groups 5 and 6 (Figure 6).

30 [0071] The ability of the antibodies raised in the immunized monkeys to neutralize HIV-1BX08 virus in human PBMC was assayed based on the reduction of p24 levels.

[0072] The neutralization assay was performed essentially as described in reference 18. Briefly, serum dilutions were mixed with HIV-1 BX08 and the mixtures incubated for 1 hour, then added to susceptible human PBMC cells. Titres were recorded as the dilution of serum at which p24 was reduced by 80%.

5 Serum samples were assayed at 1:2, 1:8 and 1:32 dilution on the virus (1:6, 1:24 and 1:26 dilutions after the addition of cells). p24 levels were evaluated by p24-specific ELISA assay.

[0073] DNA vaccination on its own, group 5, and ALVAC on its own, group 6, had no monkeys showing reduction of p24 levels greater than 80%. The
10 low DNA (600 ug) plus ALVAC, group 4, also showed no monkeys with greater than 80% reduction of p24 titres. VLP plus DNA, either high or low dose (group 1 and 2) showed enhanced reduction of p24 levels compared to VLPs alone, group 7. High dose DNA, group 3, in combination with ALVAC enhanced the ability to elicit p24 or virus neutralising antibodies over the low dose, group 4 or ALVAC
15 alone, group 6. These results indicate that DNA vaccination in combination with VLPs or ALVAC enhanced the levels of virus neutralising antibodies as indicated by the reduction of p24 levels in the sera of the immunized monkeys.

[0074] The percentage reduction of p24 is calculated relative to the amount of p24 produced in the presence of the corresponding dilution of week 2
20 samples.

SUMMARY OF DISCLOSURE

[0075] In summary of this disclosure, the present invention provides novel immunization procedures and immunogenic compositions for generating virus neutralizing levels of antibodies to a primary HIV isolate and vectors utilized
25 therein and for the generation of components for use therein. Modifications are possible within the scope of this invention.

Table 1 Study Design

| Group number | Treatment – Week 0, 4 | Treatment – Week 24,44 |
|--------------|--|--|
| 1 | 3 mg BX08 DNA 50 µg BX08 VLP 100 µg QS21 | 3 mg BX08 DNA 50 µg BX08 VLP 100 µg QS21 |
| 2 | 600 µg BX08 DNA 50 µg BX08 VLP 100 µg QS21 | 600 µg BX08 DNA 50 µg BX08 VLP 100 µg QS21 |
| 3 | 3 mg BX08 DNA | ALVAC(2) BX08 (1×10^8 pfu) |
| 4 | 600 µg BX08 DNA | ALVAC(2) BX08 (1×10^8 pfu) |
| 5 | 3 mg BX08 DNA | 3 mg BX08 DNA |
| 6 | Control DNA | ALVAC(2) BX08 (1×10^8 pfu) |
| 7 | 50 µg BX08 VLP 100 µg QS21 | 50 µg BX08 VLP 100 µg QS21 |

Table 2 Number of Monkeys showing > 80% reduction of p24 titre.

| Group number | Week 26 Bleed | Week 44 Bleed |
|--------------|---------------|---------------|
| 1 | 3/4 | 3/4 |
| 2 | 3/4 | 4/4 |
| 3 | 2/4 | 2/4 |
| 4 | 0/4 | 0/4 |
| 5 | 0/4 | 0/4 |
| 6 | 0/4 | 0/4 |
| 7 | 2/4 | 3/4 |

REFERENCES

1. UNAIDS, WHO. AIDS epidemic update: December 1999. Geneva: World Health Organisation; 1999.
2. Heyward et al 1998. HIV vaccine development and evaluation: realistic expectations. *AIDS Res Hum Retrovir* 14:S205-S210.
3. Haigwood NL and Zolla-Pazner S. 1998. Humoral immunity to HIV, SIV and SHIV. *AIDS* 12: S121-S132.
4. Johnson et al . 1998. Cellular immune responses to HIV-1. *AIDS* 12:S113-120.
5. Keefer, M. et al 1996. *AIDS Res Hum Retroviruses*. 12:683-693.
6. Cox W. et al 1993. *Virology* 195:845-850.
7. Graham BS, Keefer MC, McElrath MJ, Gorse GJ, Schwartz DH, Weinhold K, Matthews TJ, Esterlitz JR, Sinangil F, Fast PE . 1996. *Ann Intern Med* Aug 15;125(4):270-9.
8. Pincus SH, Messer KG, Cole R, Ireland R, VanCott TC, Pinter A, Schwartz DH, Graham BS, Gorse GJ. 1997. *J Immunol* 1997 Apr 1;158(7):3511-20.
9. Fang ZY et al. 1999. *J Infect Dis*. 180(4):1122-32.
10. Rovinski B et al. 1995. *AIDS Res Hum Retroviruses*. 11:1187-1195.
11. Arp J. et al. 1999. *Viral Immunolgy* 12(4):281-296.
12. Zolka-Pazner et al, *J. Virology*, vol. 72:1052-1059, 1998.
13. Karin M and Richards RI. Human metallothionein genes - primary structure of the metallothionein-II gene and a related processed gene. *Nature* 1982;299:797-802.
14. Sambrook J *et al*. *Molecular Cloning A Laboratory Manual*, Second Ed.: Cold Spring Harbour Laboratory Press. 1989.
15. Haynes JR, Cao SX, Rovinski B, Sia C, James O, Dekaban GA, Klein MH. Production of immunogenic HIV-1 viruslike particles in stably engineered monkey cell lines. *AIDS Res Hum Retroviruses* 1991;7:17-27.

16. Persson R, Cao S, Cates G, Yao F, Klein M, Rovinski B. Modifications of HIV-1 Retrovirus-like Particles to Enhance Safety and Immunogenicity. *Biologicals* 1998, 26(4): 255-265.
17. Southern PJ and Berg P. Transformation of mammalian cells to antibiotic resistance with a bacterial gene under control of the SV40 early region promoter. *J Molec Appl Genet* 1982;1: 327-341.
18. Graham, B.S. et al, 1993, *J. Infection Des.* 167:533-537.
19. Tartaglia, J. et al., 1992, *Virology*, 188:219-232.
20. Piccini, A. et al., 1987, *Methods in Enzymology* 153:545-563.

CLAIMS

What we claim is:

1. A method for generating in a host a virus neutralizing level of antibodies to a primary HIV isolate, comprising:

at least one administration of a priming antigen to the host, wherein the priming antigen comprises a DNA molecule encoding an envelope glycoprotein of a primary isolate of HIV-1,

resting the host for at least one specific resting period to provide for clonal expansion of an HIV antigen specific population of precursor B-cells therein to provide a primed host, and

at least one administration of a boosting antigen to the primed host to provide said neutralizing levels of antibodies, wherein the boosting antigen is selected from the group consisting of a non-infectious, non-replicating, immunogenic HIV-like particle having at least the envelope glycoprotein of a primary isolate of HIV-1 and an attenuated viral vector expressing at least an envelope glycoprotein of a primary isolate of HIV-1.

2. The method of claim 1 wherein said primary isolate is Bx08.

3. The method of claim 2 wherein said DNA molecule is contained in a plasmid vector under the control of a heterologous promoter for expression of the envelope glycoprotein in the host.

4. The method of claim 3 wherein the promoter is the cytomegalovirus promoter.

5. The method of claim 4 wherein the vector has the identifying characteristics of pCMV3Bx08 shown in Figure 2.

6. The method of claim 1 wherein the at least one administration of a priming antigen is at least two administrations of the priming antigen.

7. The method of claim 6 wherein the at least one specific resting period is effected after each priming administration.

8. The method of claim 1 wherein the at least one specific resting period is between about 2 months to about 12 months.

9. The method of claim 1 wherein said non-infectious, non-replicating, immunogenic HIV-like particle comprises an assembly of:

(i) an *env* gene product,

(ii) a *pol* gene product, and

(iii) a *gag* gene product,

said particle being encoded by a modified HIV genome deficient in long terminal repeats (LTRs) and containing *gag*, *pol* and *env* in their natural genomic arrangement.

10. The method of claim 9 wherein the *env* gene is that from primary isolate BX08.

11. The method of claim 1 wherein said non-infectious, non-replicating, immunogenic HIV-like particle is administered in conjunction with an adjuvant.

12. The method of claim 11 wherein the adjuvant is QS21.

13. The method of claim 1 wherein said attenuated viral vector is an attenuated avipoxvirus

14. The method of claim 13 wherein the attenuated viral vector contains a modified HIV-genome deficient in long terminal repeats, wherein at least the *env* gene is that from primary isolate BX08.

15. The method of claim 14 wherein the attenuated avipoxvirus vector is the attenuated canary poxvirus ALVAC.

16. The method of claim 15 wherein the attenuated canary poxvirus vector has the identifying characteristics of vCP1579.

17. The method of claim 1 wherein the at least one administration of a boosting antigen is at least two administrations of a boosting antigen.

18. A vector, comprising a DNA sequence encoding an envelope glycoprotein of a primary isolate of HIV-1 under the control of a heterologous promoter for expression of the envelope glycoprotein in a host organism.

19. The vector of claim 18 wherein the vector is a plasmid vector.

20. The vector of claim 18 wherein said primary HIV-1 isolate is Bx08.

21. The vector of claim 20 wherein the promoter is the cytomegalovirus promoter.

22. The vector of claim 21 which has the identifying characteristics of pCMV3Bx08 shown in Figure 2.

23. The vector of claim 18 wherein the vector is an attenuated viral vector.

24. The vector of claim 23 wherein the attenuated viral vector is a attenuated avipoxvirus vector.

25. The vector of claim 24 wherein the attenuated avipoxvirus vector is the attenuated canary poxvirus vector ALVAC.

26. The vector of claim 25 wherein the attenuated viral vector has the identifying characteristics of vCP1579 shown in Figure 4.

27. A vector, comprising a modified HIV genome deficient in long terminal repeats and a heterologous promoter operatively connected to said genome for expression of said HIV genome in mammalian cells to produce non-infectious, non-replicating and immunogenic HIV-like particles, wherein at least the *env* gene is that from a primary isolate of HIV-1.

28. The vector of claim 27 wherein the vector is a plasmid vector.

29. The vector of claim 28 wherein the primary HIV-1 isolate is BX08.

30. The vector of claim 29 wherein the promoter is type IIA metallothionein promoter.

31. The vector of claim 30 which has the identifying characteristics of p133B1 shown in Figure 3.

Figure 1 Plasmid pCMV.Bx08.gp160




Figure 2 DNA immunization plasmid pCMV3Bx08.




Figure 3. Pseudovirion Expression Plasmid p133B1 HIV-1 Bx08




FIGURE 4

ALVAC(2)120(BX08)GNP

(vCP1579)

(ALVAC *Xho*I Restriction Map)




Figure 1: Time line of PHE-vaccine 11





Figure 5 continued








Figure 2. 26 wk macaque immunoreactivity to HIV antigens (1:100 diln)



667
Figure 3. 26 wk macaque serum immunoreactivity to HIV antigens (1:1000 diln)

2-14-POX R

Fig



2-110-HIV




Figure 10 Plasmid pHIV76




Figure 11

vCP1579: H6/HIV Pol/Nef epitope cassette in ALVAC C5 site

```

1  TTTTTTTCAT TATTTAGAAA TTATGCATTT TAGATCTTTA TAAGCGGCCG TGATTAAC TA
61  GTCATAAAAA CCCGGGATCG ATTCTAGACT CGAGGGTACC GGATCTTAAT TAATTAGTCA
121 TCAGGCAGGG CGAGAACGAG ACTATCTGCT CGTTAATTAA TTAGGTCGAC GGATCCCCCA
181 ACAAAAAC TA ATCAGCTATC GGGGTTAATT AATTAGTTAT TAGACAAGGT GAAAACGAAA
241 CTATTTGTAG CTTAATTAAT TAGAGCTTCT TTATTCTATA CTTAAAAAGT GAAAATAAAT
301 ACAAAGGTTT TTGAGGGTTG TGTAAATTG AAAGCGAGAA ATAATCATAA ATTATTTT CAT
361 TATCGCGATA TCCGTTAAGT TTGTATCGTA ATGCCACTAA CAGAAGAAGC AGAGCTAGAA
421 CTGGCAGAAA ACAGAGAGAT TCTAAAAGAA CCAGTACATG GAGTGTATTA TGACCCATCA
481 AAAGACTTAA TAGCAGAAAT ACAGAAGCAG GGGCAAGGCC AATGGACATA TCAAATTTAT
541 CAAGAGCCAT TTAAAAATCT GAAAACAGGA ATGGAGTGGA GATTTGATTC TAGATTAGCA
601 TTTCATCACG TAGCTAGAGA ATTACATCCT GAATATTTTA AAAATTGTAT GGCAATATTC
661 CAAAGTAGCA TGACAAAAAT CTTAGAGCCT TTTAGAAAAC AAAATCCAGA CATAGTTATC
721 TATCAATACA TGGATGATTT GTATGTAGGA TCTGACTTAG AAATAGGGCA GCATAGAACA
781 AAAATAGAGG AGCTGAGACA ACATCTGTTG AGGTGGGGAC TTACAACCAT GGTAGGTTTT
841 CCAGTAACAC CTCAAGTACC TTAAAGACCA ATGACTTACA AAGCAGCTGT AGATCTTTCT
901 CACTTTTAA AAGAAAAAGG AGGTTTAGAA GGGCTAATTC ATTCTCAACG AAGACAAGAT
961 ATTCTTGATT TGTGGATTTA TCATACACAA GGATATTTTC CTGATTGGCA GAATTACACA
1021 CCAGGACCAG GAGTCAGATA CCCATTAACC TTTGGTTGGT GCTACAAGCT AGTACCAATG
1081 ATTGAGACTG TACCAGTAAA ATTAAAGCCA GGAATGGATG GCCCAAAGT TAAACAATGG
1141 CCATTGACAG AAGAAAAAAT AAAAGCATTA GTAGAAATTT GTACAGAGAT GGAAAAGGAA
1201 GGGAAAATTT CAAAAATTGG GCCTTAATTT TTCTGCAGCC CGGGGGATCC TTTTATAGC
1261 TAATTAGTCA CGTACCTTTG AGAGTACCAC TTCAGCTACC TCTTTTGTGT CTCAGAGTAA
1321 CTTTCTTTAA TCAATTCCAA AACAG

```

Upstream (right) flanking sequence: 1-266

VV H6 promoter: 267-390

HIV pol/nef/pol/nef/pol cassette: 391-1227

Downstream (left) flanking sequence: 1227-1345

Figure 12

E3L and K3L genes in C6

10 20 30 40 50 60 70 80 90 100 110
GAGCTCGCGG CCGCCTATCA AAAGTCTTAA TGAGTTAGGT GTAGATAGTA TAGATATTAC TACAAAGGTA TTCATATTTC CTATCAATTC TAAAGTAGAT GATATTAATA
CTCGAGCGCC GCGGATAGT TTTCAGAATT ACTCAATCCA CATCTATCAT ATCTATAATG ATGTTTCCAT AAGTATAAAG GATAGTTAAG ATTTCACTCA CTATAATTAT

120 130 140 150 160 170 180 190 200 210 220
ACTCAAAGAT GATGATAGTA GATAATAGAT ACGCTCATAT AATGACTGCA AATTGGAGG GTTCACATTT TAATCATCAC GCGTTCATAA GTTCAACTG CATAGATCAA
TGAGTTTCTA CTACTATCAT CTATTATCTA TCGAGTATA TTAGTGAGCT TTAACCTGCG CAAGTGTAAG ATTAGTAGTG CGCAAGTATT CAAAGTTGAC GTATCTAGTT

230 240 250 260 270 280 290 300 310 320 330
AATCTCACTA AAAAGATAGC CGATGTATTT GAGAGAGATT GGACATCTAA CTACGCTAAA GAAATTACAG TTATAAATAA TACATAATGG ATTTTGTAT CATCAGTTAT
TTAGAGTGAT TTTTCTATCG GCTACATAAA CTCTCTCTAA CCTGTAGATT GATCGGATTT CTTTAATGTC AATATTTTAT ATGTATTACC TAAACAATA GTAGTCAATA

340 350 360 370 380 390 400 410 420 430 440
ATTTAACATA AGTACAATAA AAAGTATTAA ATAAAATAC TTAGTTACGA AAAAATGACT AATTAGCTAT AAAAACCAG ATCTCTCGAG GTCGACGGTA TCGATAAGCT
TAAATTGAT TCTGTATT TTTTCAATTT TATTTTATG AATGAATGCT TTTTACTGA TTAATCGATA TTTTGGGTC TAGAGAGCTC CAGCTGCCAT AGCTATTCCA

450 460 470 480 490 500 510 520 530
TGATATCGAA TTCTATAAAA TT A TTG ATG TCT ACA CAT CCT TTT GTA ATT GAC ATC TAT ATA TCC TTT TGT ATA ATC AAC TCT AAT CAC TTT
ACTATAGCTT AAGTATTTT AA T AAC TAC AGA TGT GTA GGA AAA CAT TAA CTG TAG ATA TAT AGG AAA ACA TAT TAG TTG AGA TTA GTG AAA
Q H R C M R K Y N V D I Y G K T Y D V R I V K
-----K3L-----

540 550 560 570 580 590 600 610 620
AAC TTT TAC AGT TTT CCC TAC CAG TTT ATC CCT ATA TTC AAC ATA TCT ATC CAT ATG CAT CTT AAC ACT CTC TGC CAA GAT AGC TTC AGA
TTG AAA ATG TCA AAA GGG ATG GTC AAA TAG GGA TAT AAG TTG TAT AGA TAG GTA TAC GTA GAA TTG TGA GAG ACG GTT CTA TCG AAG TCT
V K V T K G V L K D R Y E V Y E D M H M K V S E A L I A E S
-----K3L-----

630 640 650 660 670 680 690 700 710
GTG AGG ATA GTC AAA AAG ATA AAT GTA TAG ACG ATA ATC CTT CTC GTA TAC TCT GCC CTT TAT TAC ATC GCC CGC ATT GGG CAA CGA ATA
CAC TCC TAT CAG TTT TTC TAT TTA CAT ATC TCG TAT TAG GAA GAG CAT ATG AGA CGG GAA ATA ATG TAG CGG GCG TAA CCC GTT GCT TAT
H P Y D F L Y I Y L A Y D K E Y V R G K I V D G A N P L S Y
-----K3L-----

720 730 740 750 760 770 780 790 800 810
ACA AAA TGC ARG CMT ACG ATACAACTT AACGGATATC GCGATAATGA AATAATTTAT GATTATTTCT CGCTTTCAAT TTAACACAAC CCTCAAGAAC
TGT TTT ACG TTC GTA TGC TATGTTGAA TTGCTATAG CGCTATTACT TTATTAAATA CTAATAAAGA CGGAAGTTA AATTGTGTTG GGAGTTCTTG
C F A L M
-----K3L-----

820 830 840 850 860 870 880 890 900 910 920
CTTTGTATTT ATTTTACCTT TTTAAGTATA GAATAAAGAA AGCTCTAATT AATTAAAGAA CAGATTGTTT CGTTTTCGCC TTGGCGTATC ACTAATTAAT TAACCGGGC
GAACATAAA TAAAGTGAA AAATTCATAT CTATTCTCTT TCGAGATTAA TTAATTACTT GTCTAACAAA GCAAAAGCGG AACCGCATAG TGATTAAATA ATTGGGCGCG

930 940 950 960 970 980 990 1000 1010 1020 1030
TGCAGCTCGA GGAATTCAC TATATCGACA TATTTCATTT GTATACACAT AACCATTAAT AACGTAGAAT GTATAGGAAG AGATGTAAAG GGAACAGGT TTGTTGATTC
ACGTGAGCT CCTTAAGTGT ATATAGCTGT ATAAAGTAAA CATATGTGTA TTGGTAATGA TTGCATCTTA CATATCTTTC TCTACATTGC CTTGTGCCA AACCACTAAG

1040 1050 1060 1070 1080 1090 1100 1110 1120 1130 1140
GCAACTATT CTAATACATA ATTCTCTCTT TAATACGTCT TGCAAGTAAT CTATTATAGA TGCCAAGATA TCTATATAAT TATTTTGTAA GATGATGTTA ACTATGTGAT
CGTTTGATAA GATTATGTAT TAAGAAGACA ATATOCAGA ACGTGCAATTA GATAATATCT ACGGTTCTAT AGATATATTA ATAAACATT CTACTACAAT TGATACACTA

1150 1160 1170 1180 1190 1200 1210 1220 1230 1240 1250
CTATATAAGT AGTGTATTA TTCTATGATT TCGATATATG TTCCAACCTT GTCTTTGTGA TGTCTAGITT CGTAAATCTT ATAGCATCTT CAAAAATAT ATTGCGATAT
GATATATCA TCACATTATT AAGTACATA AGCTATATAC AAGGTTGAGA CAGAACACT ACAGATCAAA GCATTATAGA TATCGTAGGA GTTTTTATA TAAGCGTATA

1260 1270 1280 1290 1300 1310 1320 1330 1340 1350 1360
ATTCCCAAGT CTTCACTCTT ATCTTCTAAA AAATCTTCAA CGTATGGAAT ATAATAATCT ATTTTACCTC TTCTGATATC ATTAATGATA TAGTTTTGA CACTATCTTC
TAAGGGTTCA GAAGTCAAGA TAGAAGATTT TTTAGAAGTT GCATACCTTA TATTATTAGA TAAATGGAG AAGACTATAG TAATTACTAT ATCAAAACT GTGATAGAAG

1370 1380 1390 1400 1410 1420 1430 1440 1450 1460 1470
TGTCAATTGA TTCTTATCA CTATATCTAA GAAACGGATA GCGTCCCTAG GACGAACACT TGCCATTAAAT ATCTCTATTA TAGCTTCTGG ACATAATTCA TCTATTATAC
ACAGTTAACT AAGAATAAGT GATATAGATT CTTTGCTAT CCGAGGGATC CTGCTTGATG ACGTAATTA TAGAGATAT ATCGAAGACC TGTATTAGT AGATAATATG

1480 1490 1500 1510 1520 1530 1540 1550 1560 1570 1580
CAGAATTAAT GGAACATTT CCGTATCTAT CTAACATAGT TTTAAGAAAG TCAGAACTCA AGACCTGATG TTCAATATTT GGTTCATACA TGAATGATC TCTATTGATG
GTCTTAATTA CCTTGATAA GGCATAGATA GATTGTATCA AAATCTTTC AGTCTTAGAT TCTGGACTAC AAGTATATA CCAAGTATG ACTTTACTAG AGATAACTAC

1590 1600 1610 1620 1630 1640 1650 1660 1670 1680 1690
ATAGTACTA TTTCATTCTC TGAAATTTGG TAACTCATTC TATATATGCT TTCTTGTTG ATGAAGGATA GAATATATCTC AATAGAATTT GTACCAACAA ACTGTTCTCT
TATCACTGAT AAAGTAAGAG ACTTTTARCC ATTGAGTAAG ATATATAAGA AAGGAACAACT TACTTCTAT CTTATATGAG TTATCTTAAA CATGGTGTGT TGACAAGAGA

1700 1710 1720 1730 1740 1750 1760 1770 1780 1790 1800
TATGAATCGT ATATCATCAT CTGAATAAT CATGTAAGGC ATACATTAA CAATTAGAGA CTTGCTCTCT GTTATCAATA TACTATTCTT GTGATAATTT ATGCTGAGG
ATACTTAACA TATAGTAGTA GACTTTATTA GTACATTCCG TATGTAAATT GTTAATCTCT GAACAGAGGA CAATAGTTAT ATGATAAGAA CACTATTAAA TACACACTCC

1810 1820 1830 1840 1850 1860 1870 1880 1890 1900 1910
CAAAATTTGTC CACGTTCTTT AATTTTGTTA TAGTAGATAT CAAATCCANT GGAGCTACAG TTCTTGCTT AAACAGATAT AGTTTTCTG GAACAAATTC TACAACATTA
GTTTAAACAG GTGCAAGAAA TTAACAAT ATCATCTATA GTTTAGGTTA CTTGCTCTC AAGAACCAG TTTGCTTAA TCAAAAGAC CTTGTTTAA AGTTGTAAAT

1920 1930 1940 1950 1960 1970 1980 1990 2000 2010 2020
TTATAAAGCA CTTTGGGTAG ATAAGTGGCA TGAATCCTA TTTTAATTAA TGCTATCGCA TTGCTCTCT GCAATATCC AAACGCTTT GTGATAGTAT GGCATTCAAT
AATATTTCTT GAAACCATC TATTCACCTT ACTTTAGGAT AAAATTAATT ACGATAGCGT AACAGGAGCA CGTTTATAGG TTTGCGAAA CACTATCTA CCGTAAGTAA

2030 2040 2050 2060 2070 2080 2090 2100 2110 2120 2130
GTCTAGAAC GCTCTACGAA TATCTGTGAC AGATATCATC TTTAGAGAA ATACTAGTCG CGTTAATAGT ACTACAATTT GTATTTTITA ATCTATCTCA ATAAAAAAT
CAGATCTTTC CGAGATGCTT ATAGACACTG TCTATAGTAG AAATCTCTTA TATGATCAGC GCAATTATCA TGATGTTAAA CATAAAAAAT TAGATAGAGT TATTTTITTA

2140 2150 2160 2170 2180 2190 2200 2210 2220 2230
TAATATGAT GATTCAATGT ATAACTAAC TACTAATGT TATTGATAAC TAGAATCA GAA TCT AAT GAT GAC GTA ACC AAG AAG TTT ATC TAC TGC CAA
ATTATACATA CTAAGTTACA TATTGATTTC ATGATTGACA ATAACATTG ATCTTAGT CTT AGA TTA CTA CTG CAT TGG TTC TTT AAA TAG ATG ACG GTT
-----E3L-----

2240 2250 2260 2270 2280 2290 2300 2310 2320
TTT AGC TGC ATT ATT TTT AGC ATC TCG TTT AGA TTT TCC ATC TGC CTT ATC GAA TAC TCT TCC GTC GAT GTC TAC ACA GGC ATA AAA TGT
AAA TCG ACG TAA TAA AAA TCG TAG AGC AAA TCT AAA AGG TAG ACG GAA TAG CTT ATG AGA AGG CAG CTA CAG ATG TGT CCG TAT TTT ACA
<K A A N N K A D R K S K G D A K D F V R G D I D V C A Y P T

2330 2340 2350 2360 2370 2380 2390 2400 2410
AGG AGA GTT ACT AGG CCC AAC TGA TTC AAT ACG AAA AGA CCA ATC TCT CTT AGT TAT TTG GCA GEA CTC ATT AAT AAT GGT GAC AGG GTT
TCC TCT CAA TGA TCC GGG TTG ACT AAG TTA TGC TTT TCT GGT TAG AGA GAA TCA ATA AAC CGT CAT GAG TAA TTA CCA CTC TCC CAA
<P S N S P G V S E I R F S W D R K T I Q C Y E N I I T V P N

2420 2430 2440 2450 2460 2470 2480 2490 2500
AGC ATC TTT CCA ATC AAT AAT TTT TTT AGC CCG AAT AAC ATC ATC AAA AGA CTT ATG ATC CTC TCT CAT TGA TTT TTC CCG GGA TAC ATC
TCG TAG AAA GGT TAG TTA TTA AAA AAA TCG GCC TTA TTG TAG TAG TTT TCT GAA TAC TAG GAG AGA GTA ACT AAA AAG CGC CCT ATG TAG
<A D K W D I I K K A P I V D D F S K H D E R N S K E R S V D

2510 2520 2530 2540 2550 2560 2570 2580 2590
ATC TAT TAT GAC GTC AGC CAT AGC ATC AGC ATC CCG CTT ATC CCG CTC CGT TGT CAT AAA CCA ACG AGG AGG AAT ATC GTC GGA GCT GTA
TAG ATA ATA CTG CAG TCG GTA TCG TAG TCG TAG GCC GAA TAG GCG GAG GCA ACA GTA TTT GGT TGC TCC TCC TTA TAG CAG CCT CGA CAT
<D I I V D A M A D A D P K D A E T T M P W R P P I D D S S Y

2600 2610 2620 2630 2640 2650 2660 2670 2680
CAC CAT AGC ACT ACG TTG AAG ATC GTA CAG AGC TTT ATT AAC TTC TCG CTT CTC CAT ATT AAG TTG TCT AGT TAG TTG TGC AGC AGT AGC
GTG GTA TCG TGA TGC AAC TTC TAG CAT GTC TCG AAA TAA TTG AAG AGC GAA GAG GTA TAA TTC AAC AGA TCA ATC AAC ACG TCG TCA TCG
<V M X S R Q L Y L A K N V E R K E M N L O R T L Q A A T A

2690 2700 2710 2720 2730 2740 2750 2760 2770
TCC TTC GAT TCC AAT GTT TTT AAT AGC CCG ACA CAC AAT CTC TGC GTC AGA ACG CTC GTC AAT ATA GAT CTT AGA CAT TT TTAGAGAGAA
AGG AAG CTA AGG TTA CAA AAA TTA TCG GCG TGT GTG TTA GAG ACG CAG TCT TGC GAG CAG TTA TAT CTA GAA TCT GTA AA AATCTCTCTT
<G E I G I N K I A A C V I E A D S R E D I Y I K S M

2780 2790 2800 2810 2820 2830 2840 2850 2860 2870 2880
CTAACACAAC CAGCAATAAA ACTGAACCTA CTTTATCAAT TTTTATTCA TCACTCTCTG GTGGTCTGTC GTTCTATCG AATGTAGCTC TGAATAACC GTCATCTATA
GATTGTOTTG GTGTTTATTT TGACTTGGAT GAAATAGTAA AAAAATAAGT AGTAGGAGAC CACCAAGCAG CAAGATAGC TTACATCGAG ACTAATTGGG CAGTAGATAT

2890 2900 2910 2920 2930 2940 2950 2960 2970 2980 2990
GGTATGCTG GTTCTGGAGA TTCTGGAGGA GATGATATAT TATCTGGAAG AATCTCTGTT ATTTCTTGT TTTCTATGAT CGATTGCGTT GTAAACATTAA GATTGCGAAA
CCACTACGAC CAAGACCTCT AAGACCTCTT CTACCTAATA ATAGACCTTC TTAGAGACAA TAAAGGAACA AAAGTACATA GCTAACGCAA CATTGTAAAT CTACGCTTT

3000 3010 3020 3030 3040 3050 3060 3070 3080 3090 3100
TGCTCTAAAT TTGGGAGGCT TAAAGTGTG TTTGCAATCT CTACACGGCT GTCTAAGTAG TGGAGTTGTC TCAGCTGCTC TAGTTTGAAT CATCATCGGC GTAGTATTC
ACGAGATTTA AACCTTCCA ATTTCAAC AAACGTTAGA GATGTGCGCA CAGATTGATC ACCTCCAAGC AGTCAAGCAG ATCAAACTTA GTAGTAGCGG CATCATAGG

15/15

3110 3120 3130 3140 3150 3160 3170 3180 3190 3200 3210
TACTTTTACA GTTAGGACAC GGTGTATTGT ATTTCTCGTC GAGAAGCTTA AAATAATCGT TGTAACTCAC ATCCTTTATT TTATCTATAT TGTATTCTAC TCCTTTCTTA
ATGAAAATGT CAATCCCTGT CCACATAACA TAAAGAGCAG CTCCTGCAAT TTATTAGCA ACATGAGGTG TAGGAAATAA AATAGATATA ACATAAGATG AGGAAAGAAT

3220 3230 3240 3250 3260 3270 3280 3290 3300 3310 3320
ATGCAITTTA TACCGAATAA GAGATAGCGA AGGAATTCIT TTTATIGATT AACTAGTCAA ATGAGTATAT ATAAATGAAA AAGTAAATA TAAATCATAT AATAATGAAA
TACGTAAAT ATGGCTTATT CTCTATCGCT TCCTTAAGAA AAATAACTAA TTGATCAGTT TACTCATATA TATTAACTTT TTCATTTTAT ATTTAGTATA TTATTACTTT

3330 3340 3350 3360 3370 3380 3390 3400 3410 3420 3430
CGAATATCA GTAATAGACA GGAAGTGGCA GATTCTTCTT CTAATGAAGT AAGTACTGCT AAATCTCCAA AATTAGATAA AATGATACA GCAATACAG CTCATTCAA
GCTTTATAGT CATTATCTGT CCTTGACCGT CTAAGAAGAA GATTACTTCA TTCATGACGA TTTAGAGGTT TTAATCTATT TTTACTATGT CGTTTATGTC GAACTAAGTT

3440 3450 3460 3470 3480 3490 3500 3510 3520 3530 3540
CGAATTACCT TTTAATTTT TCAGACACAC CTTATTACAA ACTAACTAAG TCAGATGATG AGAAAGTAAA TATAAATTTA ACTTATGGGT ATAATATAAT AAAGATTCAT
GCTTAATGGA AAATTAAAA AGTCTGTGTG GAATAATGTT TGATTGATTC AGTCTACTAC TCTTTCATTT ATATTTAAAT TGAATACCCA TATTATATTA TTTCTAAGTA

3550 3560 3570 3580 3590 3600 3610 3620 3630 3640 3650
GATATTAAAT ATTTACTTAA CGATGTIAAT AGACTTATTC CATCAACCCC TTCAAACCTT TCTGGATATT ATAAAAATACC AGTTAATGAT ATTAAAAATAG ATTGTTTAAAG
CTATAATAT TAAATGAAT GCTACAATTA TCTGAATAAG GTAGTTGGCG AAGTTTGGAA AGACCTATTA TATTTTATGG TCAATTACTA TAATTTTATC TAACAAATTC

3660 3670 3680 3690 3700 3710 3720 3730 3740 3750 3760
AGATGTAAAT AATTATTTGG AGGTAAAGCA TATAAAATTA GTCTATCTTT CACATGGAAA TGAATTACCT AATATTAAAT ATTATGATAG GAATTTTITA GGATTTACAG
TCTACATTTA TTAATAAACC TCCATTTCCT ATATTTTAAAT CAGATAGAAA GTGTACCTTT ACTTAATGGA TTATAATTAT TAATACTATC CTTAAAAAAT CCTAAATGTC

3770 3780 3790 3800 3810 3820 3830 3840 3850 3860 3870
CTGTATATG TATCAACAAT ACAGGCAGAT CTATGTTTAT GGTAAAACAC TGTACGGGA AGCAGCATTC TATGGTAACT GGCCTATGTT TAATAGCCAG ATCATTTTAC
GACAATATAC ATAGTGTGTA TGTCCGTCTA GATACCAATA CCATTTTGTG ACATTTGCCCT TCGTGGTAAG ATACCAATGA CCGGATACAA ATTATCGGTC TAGTAAATG

3880 3890 3900 3910 3920 3930 3940 3950 3960 3970 3980
TCTATAACA TTTTACCACA AATAATAGGA TCCTCTAGAT ATTTAATATT ATATCTAACA ACAACAAAA AATTAAAGGA TGTATGGCCA GAATATTTT CTACTAATAA
AGATATTGT AAAATGGTGT TTATTATCCT AGGAGATCTA TAAATTATAA TATAGATTGT TGTGTTTTT TTAATTTGCT ACATACCGGT CTTCAATAAA GATGATTATT

3990 4000 4010 4020 4030 4040 4050 4060 4070 4080 4090
AGATAAGAT AGTCTATCTT ATCTACAAGA TATGAAAGAA GATAATCATT TAGTAGTAGC TACTAATATG GAAAGAAATG TATACAAAA CGTGGAAAGCT TTTATATTA
TCTATTTCTA TCAGATAGAA TAGATGTTCT ATACTTTCTT CTATTAGTAA ATCATCATCG ATGATTATAC CTTTCTTTAC ATATGTTTTT GCACCTTCGA AATATAAAT

4100 4110 4120 4130 4140 4150 4160 4170 4180 4190 4200
ATAGCATATT ACTAGAAGAT TTAATATCTA GACTTAGTAT AACAAACAG TTAATGCCA ATATCGATTC TATATTTTAT CATAACAGTA GTACATTAAAT CAGTGATATA
TATCGTATAA TGATCTTCTA AATTTTAGAT CTGATCATA TTGTTTTGTC AATTACGGT TATAGCTAAG ATATAAAGTA GTATTGTCAT CATGTAATTA GTCATATAT

4210 4220 4230 4240 4250 4260 4270 4280 4290 4300 4310
CTGAACGAT CTACAGACTC AACTATGCAA GGATAAGCA ATATGCCAAT TATGTCTAAT ATTTTAACTT TAGAACTAAA ACGTTCTACC AATCTAAAA ATAGGATACG
GACTTTGCTA GATGTCTGAG TTGATAAGTT CCTTATTCGT TATACGGTAA ATACAGATTA TAAATTTGAA ATCTTGATTT TGCAAGATGG TTATGATTTT TATCCTATGC

4320 4330 4340 4350 4360 4370 4380 4390 4400 4410 4420
TGATAGGCTG TTAAGAGCTG CAATAAATAG TAAGGATGTA GAAGAAATAC TTTGTCTAT ACCTTCGGAG GAAAGAACTT TAGAACRACT TAAGTTTAAAT CAACTTGTGA
ACTATCCGAC AATTTTCGAC GTTATTTATC ATTCCTACAT CTTCTTTATG AAACAAGATA TGGAAAGCTC CTTTCTTGAA ATCTTGTTGA ATTCAAATTA GTTTGAACAT

4430
TTTATGAAGG TACC
AAATACCTCC ATGG

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
8 November 2001 (08.11.2001)

PCT

(10) International Publication Number
WO 01/82962 A3

(51) International Patent Classification⁷: **A61K 39/21**,
39/39, C12N 15/86, C07K 14/16

(74) Agent: STEWART, Michael, L; Sim & McBurney, 6th
Floor, 330 University Avenue, Toronto, Ontario M5G 1R7
(CA).

(21) International Application Number: PCT/CA01/00577

(22) International Filing Date: 25 April 2001 (25.04.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/200,011 27 April 2000 (27.04.2000) US

(71) Applicant (*for all designated States except US*): AVEN-
TIS PASTEUR LIMITED [CA/CA]; 1755 Steeles Av-
enue West, Toronto, Ontario M2R 3T4 (CA).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM,
HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK,
LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX,
MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL,
TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): ROVINSKI, Ben-
jamin [CA/CA]; 70 Winding Lane, Thornhill, Ontario L4J
5H6 (CA). TARTAGLIA, James [US/US]; 7 Christina
Drive, Schenectady, NY 12303 (US). CAO, Shi-Xian
[CA/CA]; Apartment 408, 716 The West Mall, Etobicoke,
Ontario M9C 4X6 (CA). PERSSON, Roy [CA/CA]; Unit
604, 7 Bishop Avenue, Toronto, Ontario M2M 4J4 (CA).
KLEIN, Michel, H. [CA/CA]; 54 Strathgowen Avenue,
Toronto, Ontario M4N 1B9 (CA).

Published:

— with international search report

(88) Date of publication of the international search report:
21 March 2002

*For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*



WO 01/82962 A3

(54) Title: IMMUNIZING AGAINST HIV INFECTION

(57) Abstract: A virus neutralizing level of antibodies to a primary HIV isolate is generated in a host by a prime-boost administration of antigens. The primary antigen is a DNA molecule encoding an envelop glycoprotein of a primary isolate of HIV-1 while the boosting antigen is either a non-infectious, non-replicating HIV-like particle having the envelope glycoprotein of a primary isolate of HIV-1 or an attenuated viral vector expressing an envelope glycoprotein of a primary isolate of HIV-1.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/CA 01/00577

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A61K39/21 A61K39/39 C12N15/86 C07K14/16

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K C07K C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, BIOSIS

C. DOCUMENTS CONSIDERED TO BE RELEVANT

| Category * | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
|------------|---|-----------------------|
| X | CAVER T E ET AL: "A novel vaccine regimen utilizing DNA, vaccinia virus and protein immunizations for HIV-1 envelope presentation" VACCINE, BUTTERWORTH SCIENTIFIC. GUILDFORD, GB, vol. 17, no. 11-12, March 1999 (1999-03), pages 1567-1572, XP004158286 ISSN: 0264-410X the whole document. | 1,8,18, 19 |
| Y | | 2-7 |
| A | | 9-17 |

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *G* document member of the same patent family

Date of the actual completion of the international search

13 December 2001

Date of mailing of the international search report

02/01/2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Galli, I

INTERNATIONAL SEARCH REPORT

International Application No

PCT/CA 01/00577

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

| Category | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
|----------|--|-----------------------|
| X | WO 99 31250 A (ROVINSKI BENJAMIN ;CONNAUGHT LAB (CA); CAO SHI XIAN (CA); YAO FEI) 24 June 1999 (1999-06-24) cited in the application the whole document | 18-22, 27-29 |
| Y | see p. 12 line 2: BX08 --- | 2-7 |
| X | EVANS T.G. ET AL.: "A Canarypox vaccine expressing multiple human immunodeficiency virus type 1 genes given alone or with rgp120 elicits broad and durable CD8+ cytotoxic T lymphocyte responses in seronegative volunteers." J. INFECT. DISEASES, vol. 180, 1999, pages 290-298, XP002185651 abstract | 23-25 |
| A | --- | 26 |
| X | WO 91 05864 A (CONNAUGHT LAB) 2 May 1991 (1991-05-02) cited in the application abstract | 27,28 |
| A | figure 10 --- | 30,31 |
| P,X | WO 00 50604 A (ROVINSKI BENJAMIN ;CONNAUGHT LAB (CA); PERSSON ROY (CA); CAO SHI X) 31 August 2000 (2000-08-31) abstract | 18-22 |
| A | WO 98 40501 A (VIROGENETICS CORP) 17 September 1998 (1998-09-17) abstract ----- | 26 |

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No.

PC1/CA 01/00577

| Patent document cited in search report | | Publication date | | Patent family member(s) | Publication date |
|---|---|---------------------|----|----------------------------|---------------------|
| WO 9931250 | A | 24-06-1999 | US | 6121021 A | 19-09-2000 |
| | | | AU | 1657799 A | 05-07-1999 |
| | | | WO | 9931250 A2 | 24-06-1999 |
| | | | EP | 1038001 A2 | 27-09-2000 |
| WO 9105864 | A | 02-05-1991 | CA | 2067333 A1 | 14-04-1991 |
| | | | WO | 9105864 A1 | 02-05-1991 |
| | | | EP | 0495811 A1 | 29-07-1992 |
| | | | JP | 2537570 B2 | 25-09-1996 |
| | | | JP | 4506301 T | 05-11-1992 |
| | | | US | 5571712 A | 05-11-1996 |
| | | | US | 6291227 B1 | 18-09-2001 |
| | | | US | 5985641 A | 16-11-1999 |
| | | | US | 5439809 A | 08-08-1995 |
| WO 0050604 | A | 31-08-2000 | AU | 2789200 A | 14-09-2000 |
| | | | WO | 0050604 A1 | 31-08-2000 |
| | | | EP | 1157115 A1 | 28-11-2001 |
| WO 9840501 | A | 17-09-1998 | US | 5990091 A | 23-11-1999 |
| | | | AU | 6158498 A | 29-09-1998 |
| | | | EP | 0970226 A1 | 12-01-2000 |
| | | | JP | 2001514518 T | 11-09-2001 |
| | | | US | 6130066 A | 10-10-2000 |
| | | | WO | 9840501 A1 | 17-09-1998 |